

# **Voltammetry determination of dopamine by the electrocatalytic response of an electrode modified by a polyaniline film with an inclusion of copper(II) tetrasulfophthalocyanine**

Shaidarova L., Gedmina A., Artamonova M., Chelnokova I., Budnikov H.  
*Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

---

## **Abstract**

A polyaniline composite film with an inclusion of copper(II) tetrasulfophthalocyanine, applied on a glassy-carbon electrode (GC) has demonstrated a mediatory activity in the oxidation of dopamine in acidic and neutral media. The conditions for the formation of a polymer film on the surface of a glassy-carbon electrode for registering the maximum electrocatalytic effect in the oxidation of dopamine are found. A method of voltammetric and amperometric detection of dopamine on the composite film electrode under static and flow conditions is proposed. The use of a polyaniline composite with an inclusion of copper(II) tetrasulfophthalocyanine, coated with a Nafion film, made possible the reduction of the detection limit for dopamine to  $1 \times 10^{-8}$  M under static conditions and to 5.0 nmol under the conditions of flow-injection analysis. © 2013 Pleiades Publishing, Ltd.

<http://dx.doi.org/10.1134/S1061934813060129>

---

## **Keywords**

chemically modified electrodes, copper(II) phthalocyanine, electrocatalysis, electrooxidation of dopamine, flow-injection analysis, polyaniline